arXiv:1503.02967v2 [cond-mat.mtrl-sci] 12 Jul 2016

Critical analysis of the slope method for estimation of ice-water interfacial energy from ice nucleation experimental data

Tomáš Němec[a,b,*]

[a]Institute of Thermomechanics of the CAS, v. v. i., Dolejškova 5, 18200 Praha 8, Czech Republic
[b]New Technologies - Research Centre, University of West Bohemia, Univerzitní 8, 30614 Plzeň, Czech Republic

---

Abstract

An established procedure for the estimation of ice-water interfacial energy based on evaluation of the slope of the experimental ice nucleation rate data versus scaled temperature is critically analyzed in this work. An inconsistent estimate of the ice-water interfacial energy is found in the work of Murray *et al.* [Phys. Chem. Chem. Phys., 2010, 12, 10380–10387]. The source of the inconsistency is identified in an inappropriate regression method used for experimental ice nucleation data fitting, a correct estimate of the ice-water interfacial energy is presented, and limits of the slope method are discussed.

---

1. Introduction

The classical nucleation theory (CNT) provides a theoretical link between the ice nucleation rates and the thermophysical properties of the ice-water system, i.e. the density of the solid phase, the diffusion coefficient of the liquid phase, the equilibrium pressure, and the interfacial energy [6]. Therefore, in cases where experimental measurements of the nucleation rate are available, CNT presents an indirect, theoretical approach for the estimation of the interfacial energy provided that the other above mentioned thermophysical properties are known.

For ice and water, the interfacial energy measurements reported in literature are restricted to the triple point temperature [3]. The estimation of the interfacial energy for lower temperatures in the supercooled region, was possible only due to ice nucleation rate measurements [11] or molecular simulations of ice nuclei formation [8, 2].

In this work, a method for the estimation of ice-water interfacial energy from the experimental nucleation rates [14, 15] will be investigated that is based on the evaluation of the slope of the logarithm of the measured nucleation rate $\ln J$ vs. the function $T^{-3}(\ln(S))^{-2}$ of the experimental temperature T, where S is the ratio of water and ice saturation pressures. It will be referred to as the *slope method* in the following text. Several authors estimated the ice-water interfacial energies based on the slope method [11, 12, 9, 1] covering the temperature range 200 − 240 K. I will study the validity of the ice-water interfacial energy estimates by the slope method in this paper. Particularly, I intend to show that the ice-water interfacial energy estimate of Murray *et al.* [11] suffers from internal inconsistency and needs to be reconsidered. The nature of this inconsistency will be studied in detail to point out the weak spots of the slope method.

The paper is arranged in the following way. First, the equations of the slope method for interfacial energy estimation are derived in section 2. In section 3, a proof is presented showing the inconsistency of the interfacial energy estimate by Murray *et al.* [11]. In section 4, the source of the inconsistency is identified, a correct estimate of the ice-water interfacial energy is presented, and the limits of applicability of the slope method are discussed.

2. Slope method

The slope method [14, 15] for the estimation of the ice-water interfacial energy is based on the CNT nucleation rate equation [7]

$$J = J_0 \exp\left(-\frac{W^\star}{kT}\right) \tag{1}$$

---

[*] Corresponding author
Email address: `nemec@it.cas.cz` (Tomáš Němec)

relating the nucleation rate J [$\mathrm{m}^{-3}\mathrm{s}^{-1}$] to the nucleation work of the critical cluster $W^\star$ [J]. The pre-factor J_0 [$\mathrm{m}^{-3}\mathrm{s}^{-1}$] reflects the kinetics of the cluster growth and T [K] is absolute temperature. By using the CNT expression for the critical nucleation work $W^\star = \frac{16\pi\gamma^3 v^2}{3(kT\ln S)^2}$, Eq. (1) can be rearranged to

$$\ln J = \ln J_0 - \frac{16\pi\gamma^3 v^2}{3k^3T^3(\ln S)^2} \tag{2}$$

where γ [$\mathrm{J/m}^2$] is the ice-water interfacial energy, v [m^3] is the molecular volume of the solid phase, and the ratio of saturation pressures $S = p_l^{eq}/p_s^{eq}$ represents the supersaturation of the liquid. Here, the vapor-liquid equilibrium pressure is denoted p_l^{eq}, and the vapor-solid equilibrium pressure is denoted p_s^{eq}. The vapor-solid equilibrium pressure of cubic ice [17] is used in this work to retain consistency with the work of Murray *et al.* [11]. However, the structure of the clusters in ice nucleation is still under debate [10].

To transform Eq. (2) into the relations of the slope method, the following assumptions are made. The interfacial energy is taken constant, i.e. independent of temperature. The density of the solid phase is taken constant. And the pre-factor is taken constant. Since the size of the temperature interval of the experimental nucleation data analyzed by the slope method is typically a few Kelvins, the assumptions of the constancy of the thermophysical properties over this narrow temperature interval are plausible, and Eq. (2) can be written in the form

$$\ln J = n + mt_s \tag{3}$$

where the parameters

$$n = \ln J_0 \tag{4}$$

and

$$m = -\frac{16\pi\gamma^3 v^2}{3k^3} \tag{5}$$

are both due to the above assumptions constant, and

$$t_s = \frac{1}{T^3(\ln S)^2} \tag{6}$$

is a scaled temperature. As a result, the slope method is simply a fit of the experimental ice nucleation rate data to the linear function (3) in t_s, giving the parameters n and m. The ice-water interfacial energy γ corresponding to the analyzed ice nucleation data is recovered from the slope m according to Eq. (5), and the pre-factor J_0 is given by the absolute value n according to Eq. (4).



Figure 1: Experimental ice nucleation rates reported by Murray *et al.* [11] and Stan *et al.* [18]. The full lines show fits of the experimental data, and the dashed lines delimit the standard deviation bands. The fit of Murray *et al.* [11] is used as presented in their Fig. 4. The data of Stan *et al.* [18] as presented in their Fig. 8 are shown with the reported ±0.4 °C standard deviation.

3. Inconsistency of interfacial energy estimates

The validity of the slope method estimates of the ice-water interfacial energy based on the experimental nucleation data as published by Murray *et al.* [11] will be investigated in this section.

I have identified inconsistent estimates of interfacial energy for two specific sets of ice nucleation data investigated by Murray *et al.* [11], i.e. their own experimental data and the data measured by Stan *et al.* [18], respectively. The two experimental data sets are plotted in Fig. 1. By using the slope method, Murray *et al.* [11] estimated ice-water interfacial energy $\gamma_M = 20.8 \pm 1.2\ \mathrm{mJ/m}^2$ from their own experimental data, and ice-water interfacial energy $\gamma_S = 23.7 \pm 1.1\ \mathrm{mJ/m}^2$ from the experimental data of Stan *et al.* [18], respectively. It is the gap between the two intervals, γ_M and γ_S, that renders the two interfacial energy estimates inconsistent. In other words, as I will show below in detail, since the initial two experimental datasets used as input for the evaluation of the respective interfacial energies do overlap to a large extent, which is seen in Fig. 1, it is impossible to calculate two interfacial energy estimates by the slope method that form two disjunct intervals, as is the case with γ_M and γ_S.

To prove the particular inconsistency in interfacial energy estimates γ_M and γ_S suggested above, I will frame general requirements for a physically-relevant deduction of the ice-water interfacial energy from the nucleation rate according to the CNT. Mathematically, the deductions will utilize elementary properties of continuous, strictly monotonic functions in a reductio ad absurdum type of logical proof. The slope method analysis will present a limiting case of this general scenario.

Let us denote g a continuous, strictly monotonic function of temperature g: $T \rightarrow \gamma$, which stands for the functional dependence of the interfacial energy on temperature. Similarly, let j be a continuous, strictly monotonic function of temperature j: $T \rightarrow J$, which stands for the functional dependence of the nucleation rate on temperature, e.g. in the form of Eq. (1). Let the temperature interval $[T_1,\, T_2]$ be a subdomain of the two functions g and j that corresponds to the temperature range of the analyzed ice nucleation experiment. Both functions g and j are bijective by their above definitions. In other words, they present a one-to-one correspondence between the elements of subdomain $[T_1,\, T_2]$ and the elements of functional images $g([T_1, T_2])$ and $j([T_1, T_2])$, respectively. Therefore, inverse function j^{-1}: $J \rightarrow T$ exists and it is bijective as well. Furthermore, composite function $g_j = g(j^{-1})$: $J \rightarrow \gamma$ is bijective. Here, g_j is the formal representation of a general theoretical method to deduce the interfacial energy from the nucleation rate in temperature range $[T_1,\, T_2]$.

Now, let $\boldsymbol{J_1} = [J_{11},\, J_{12}]$ and $\boldsymbol{J_2} = [J_{21},\, J_{22}]$ be two intervals of nucleation rates that possess a non-empty intersection $\boldsymbol{J_0} = \boldsymbol{J_1} \cap \boldsymbol{J_2} \neq \emptyset$. Let $\boldsymbol{\gamma_1} = [\gamma_{11},\, \gamma_{12}]$ and $\boldsymbol{\gamma_2} = [\gamma_{21},\, \gamma_{22}]$ be g_j's functional images of intervals $\boldsymbol{J_1}$ and $\boldsymbol{J_2}$, i.e. $\boldsymbol{\gamma_1} = g_j(\boldsymbol{J_1})$ and $\boldsymbol{\gamma_2} = g_j(\boldsymbol{J_2})$, respectively. Intervals $\boldsymbol{\gamma_1}$ and $\boldsymbol{\gamma_2}$ represent the ranges of interfacial energies deduced from the two nucleation rate data sets $\boldsymbol{J_1}$ and $\boldsymbol{J_2}$ according to a general method g_j. Then $\forall J \in \boldsymbol{J_0}$ it holds that $g_j(J) \in \boldsymbol{\gamma_1}$ and $g_j(J) \in \boldsymbol{\gamma_2}$. In other words, for any value of the nucleation rate J, which belongs to the intersection of intervals $\boldsymbol{J_1}$ and $\boldsymbol{J_2}$, the deduced value of interfacial energy according to method g_j belongs to interval $\boldsymbol{\gamma_1}$ and also to interval $\boldsymbol{\gamma_2}$. Therefore, since element $\gamma = g_j(J)$ belongs to both intervals $\boldsymbol{\gamma_1}$ and $\boldsymbol{\gamma_2}$, intersection $\boldsymbol{\gamma_0} = \boldsymbol{\gamma_1} \cap \boldsymbol{\gamma_2}$ has at least one element γ and, therefore, $\boldsymbol{\gamma_0}$ must be non-empty, i.e. $\boldsymbol{\gamma_0} \neq \emptyset$.

For the particular case of the ice nucleation experimental data of Murray *et al.* [11] and Stan *et al.* [18] the intersection of the measured nucleation rate data sets is not empty, as can be seen in Fig. 1; the measured values of the nucleation rate in the two datasets clearly overlap to a large extent. According to the above reasoning the range of interfacial energies deduced from the Murray *et al.* [11] nucleation data must have an non-empty intersection with the range of interfacial energies deduced from the Stan *et al.* [18] nucleation data provided that the method used to deduce the interfacial energy assures that the temperature dependency of the interfacial energy and the nucleation rate is strictly monotonic. The strict monotonicity of both the nucleation rate and the interfacial energy is a natural requirement conforming to the physical reality; the nucleation rate decreases with increasing temperature and the interfacial energy increases with increasing temperature [6].

However, the slope method assumes, rather unphysically, that the interfacial energy is constant, and not strictly monotonic as supposed in the general case above. Under such assumption, the considerations presented above result into an even stronger requirement on the deduced interfacial energies from two intersecting nucleation rate datasets. Any constant function can be viewed as a limit of a sequence of strictly monotonic functions, e.g. a sequence of linear functions $f_n = C + a_n(T - T_0)$ with slope a_n decreasing as $1/n$ for $n \rightarrow \infty$. For every element of the sequence the above reasoning applies and the deduced interfacial energy ranges must possess a non-empty intersection. In the limit of the constant function the intersection shrinks to a single value of interfacial energy, common to both datasets. As a result, the slope method must produce interfacial energies equal to each other within their standard deviation ranges from two overlapping datasets of nucleation rates. However, we already stated that the estimates of the interfacial energy of the slope method, γ_M and γ_S, as published by Murray *et al.* [11] are not equal within their standard deviations, which is in contradiction with the outcome of the above reasoning. Therefore, the slope method procedure as preformed by Murray *et al.* [11] must contain a hidden inconsistency, an erroneous step that introduces additional uncertainty. Its nature will be discussed in the following section.

4. Discussion

The fit of the experimental ice nucleation-rate data to the linear function (3) forms the basis of



Figure 2: Comparison of the ordinary least squares regression method (dashed line) and the orthogonal distance regression method (full line) for the estimation of the slope of the ice nucleation data of Murray *et al.* [11].

the slope method. The fit results in two parameters n and m that will be discussed in this section. First, the evaluation of the slope m resulting in the estimation of ice-water interfacial energy (5) will be investigated with the goal of resolving the inconsistency found in Sec. 3. Second, the evaluation of the nucleation rate pre-factor from the term n according to Eq. (4) will be analyzed.

4.1. Interfacial energy

Although Murray *et al.* [11] do not specify the particular regression method that they used to perform the linear fit to their data yielding their reported slope $m_{OLS} = -\,(6.02 \pm 0.36)\times 10^7$ K^3, their result can be reproduced with the widely used ordinary least squares (OLS) method. By using the OLS method implemented in Python (package `scipy.stats.linregress`) to fit the Murray *et al.* [11] experimental data to the function (3) we can check that the above mentioned value of m_{OLS} is recovered. Therefore, let us assume that the OLS method was indeed used by Murray *et al.* [11] in their analysis.

The OLS method was derived with the assumption that the observation of the independent variable, i.e. the temperature in this case, is error-free. From this point of view the applicability of the OLS method is highly questionable for nucleation-rate experimental data, because the error in the temperature measurement of an ice nucleation experiment is typically larger than ±0.4 °C, and it is identified as the main source of uncertainty in the nucleation measurement [18].

The regression method derived with the assumption of non-zero observational errors in the independent variable is the orthogonal distance regression method (ODR), also known as errors-in-variables modeling, or total least squares [5]. The Python ODR implementation (`scipy.odr.odrpack`) applied to Murray *et al.* [11] ice nucleation data results in the slope $m_{ODR} = -\,(8.04 \pm 0.48)\times 10^7$ K^3.

The difference in the slope estimates between the OLS and ODR methods for the Murray *et al.* [11] ice nucleation data is shown in Fig. 2. The ice-water interfacial energy corresponding to the m_{ODR} is $\gamma_{ODR} = 22.9 \pm 0.5$ mJ/m^2 according to the slope method, Eq. (5). And after including the uncertainty in the cubic ice sublimation pressure [17] ($\pm$ 0.8 mJ/m^2), the final estimate of ice-water interfacial energy using the ODR method is $\gamma_{ODR} = 22.9$ $\pm$ 1.3 mJ/m^2. This new ODR estimate is roughly 10 % higher that the OLS estimate $\gamma_{OLS} = 20.8 \pm$ 1.2 mJ/m^2 calculated from m_{OLS}. The new value γ_{ODR} is sufficiently close to the value γ_S Murray *et al.* [11] estimated from the nucleation data of Stan *et al.* [18], and therefore γ_{ODR} does not suffer from the contradiction described in Sec. 3.

Note that the Stan data [18] are fitted correctly even with the OLS method that fails for fitting the Murray data [11]. We can check that the OLS method gives the slope $m = -8.952\times 10^7$ K^3 and the ODR method results in $m = -8.967\times 10^7$ K^3, which both correspond to the interfacial energy γ_S = 23.7 mJ/m^2 reported by Murray *et al.* [11] for the nucleation data of Stan *et al.* [18]. The Stan data, as presented in Fig. 1 (and in their Fig. 8 [18]), on the contrary to the raw data by Murray *et al.* [11], are already averaged from a set of more than 37 thousand freezing experiments, effectively removing the large uncertainty in experimental temperature measurement. The Stan data are therefore suitable for fitting even by the OLS method.

4.2. Nucleation rate pre-factor

The slope method concurrently fits two parameters, n and m, from a given data set of experimental nucleation rates. The two parameters are therefore coupled and we can expect that the larger an error in the estimation of m we make, the larger the corresponding error in n we get. It was shown in the preceding section that the interfacial energy estimate γ_M suffered from an error due to the inap-



Figure 3: Pre-factor J_0 estimated by Murray *et al.* [11] with the slope method (as given in their Table 2) from the ice nucleation data available in literature (red circles). Horizontal errorbars show the temperature range of the input experimental nucleation-rate datasets. The vertical errorbars denote the uncertainty of J_0 fit. The green full line shows the pre-factor according to the volume-diffusion controlled growth model [7]. The blue square shows the OLS-based fit of the pre-factor, and the the green diamond shows the ODR-based fit of the pre-factor from the experimental data of Murray *et al.* [11].

propriate statistical processing of the experimental data. Therefore, an error was introduced in the corresponding estimate of nucleation rate pre-factor J_0 as well.

The difference between the pre-factor estimates using the OLS and the ODR regression methods is shown in Fig. 3 for the experimental data of Murray *et al.* [11]. The fitted parameters are $n_{OLS} = 75 \pm 5$, and $n_{ODR} = 94.5 \pm 5$, respectively. By evaluating the pre-factor according to Eq. (4), we find that the OLS-based estimate of J_0 is roughly 9 orders of magnitude lower than the ODR-based result. It is the ODR-based estimate of J_0 that is within its uncertainty equal to an independent model of J_0, i.e. the volume-diffusion based description of the cluster growth [7] (p. 141), as shown in Fig. 3.

4.3. Final remarks

Not only the already discussed regression issue is the source of uncertainty in the slope method. The assumptions of the method itself, as summarized in Sec. 2, play a role as well. The constancy of the physical properties assumed by the slope method is plausible for small temperature ranges of the analyzed experimental nucleation rates. As the temperature range of the experiment increases, the interfacial energy, ice density, and the pre-factor are getting less accurately approximated by a constant. It is therefore desirable for the slope method to analyze temperature ranges as small as possible. On the other hand, to evaluate the slope of the ice nucleation data from the cloud of scattered experimental nucleation rates one needs as large temperature ranges as possible, because the uncertainty of the fit of the slope gets larger as the temperature range of the analyzed data decreases. Obviously, both requirements, i.e. having the temperature interval small for the assumptions of the slope method to be valid and having the temperature interval large for a precise evaluation of the slope, cannot be simultaneously satisfied. And uncertainty due to this fact will be inevitably present in the results of the slope method.

Fig. 3 shows the J_0 estimates from all the ice nucleation-rate datasets in the temperature range 235 – 240 K available in literature as calculated by Murray *et al.* [11]. A huge scatter in the pre-factor estimates spanning roughly 25 orders of magnitude can be observed. However, the pre-factor is only slightly temperature dependent in the case of nucleation in liquids [7] (p. 199). According to the above-mentioned diffusion-based growth model the pre-factor depends linearly on the self-diffusion coefficient of supercooled water, which was measured down to 238 K with accuracy less than 1% [16]. Therefore, the volume-diffusion based growth model gives us an estimate of the pre-factor J_0 with negligible uncertainty compared to the uncertainty of 25 orders of magnitude reported in Murray *et al.* [11]. Obviously, the scatter of J_0 estimates as derived by Murray *et al.* [11] and shown in Fig. 3 is non-physical, and reflects the errors in implementing the slope method.

Finally, the slope method was also used by Manka *et al.* [9] to estimate the ice-water energy of 15.6 mJ/m^2 from their ice nucleation data measured in the temperature range 202 – 215 K. The authors do not report any uncertainty of their interfacial energy estimate; they just state "we analyzed all of our data using the same formalism" as Murray *et al.* [11]. Therefore, a similar regression issue as in the above-discussed case of Murray *et al.* [11] interfacial energy estimate is likely to arise. Also, the relatively large temperature interval of the Manka data collides with the assumptions of constancy of the thermophysical properties of the slope method. But most importantly, the CNT formulation used in the slope method as presented in Sec. 2 is not valid for ice nucleation data measured at such low temperatures due to the omitted pressure effect, as shown recently [13]. The pressure-related terms in the CNT formulation present a considerable contribution to the nucleation work and, therefore, account for a shift in the CNT-predicted ice nucleation rate at low temperatures. The errors in CNT formulation are then projected to errors in the ice-water interfacial energy estimates of the slope method.

5. Conclusion

A method for the estimation of the ice-water interfacial energy based on the classical nucleation theory and on the evaluation of the slope of the experimental ice nucleation rates vs. a scaled temperature was thoroughly analyzed in this work. A logical contradiction in the interfacial energy estimates was identified in the work of Murray *et al.* [11] that is related to the linear regression algorithm utilized for the evaluation of the slope of experimental ice nucleation rate data. The contradiction was removed by using the orthogonal distance regression method for a proper evaluation of the slope instead of the ordinary least squares method. The corrected estimate of the interfacial energy is by 10 % (2.1 mJ/m^2) higher than the original value reported by Murray *et al.* [11].

In the light of the findings of this work, estimation methods [4, 13] utilizing the absolute values of the experimental nucleation rates instead of the temperature derivative, and using a theoretical model of the nucleation rate pre-factor instead of its fit from experimental nucleation data, present a safer way of deducing the interfacial energy from nucleation rate data avoiding uncertainties inherently contained in the slope method.

Acknowledgement

The author would like to acknowledge the institutional support RVO:61388998 of the Institute of Thermomechanics, v.v.i., support by the Czech Science Foundation (project GAP101/10/1819), by the CENTEM project (reg. no. CZ.1.05/2.1.00/03.0088) co-funded by the ERDF as part of the Ministry of Education, Youth and Sports OP RDI programme and, in the follow-up sustainability stage, supported through CENTEM PLUS (LO1402) by financial means from the Ministry of Education, Youth and Sports under the National Sustainability Programme I, and by the POLYMEM project (reg. no. CZ.1.07/2.3.00/20.0107) co-funded by the ESF as part of the Ministry of Education, Youth and Sports ECOP programme.

References

[1] A. Bhabhe, H. Pathak, and B. E. Wyslouzil. Freezing of heavy water (D2O) nanodroplets. *J. Phys. Chem. A*, 117(26):5472–5482, 2013.

[2] J. R. Espinosa, C. Vega, C. Valeriani, and E. Sanz. Seeding approach to crystal nucleation. *J. Chem. Phys.*, 144(3), 2016.

[3] L. Granasy, T. Pusztai, and P. F. James. Interfacial properties deduced from nucleation experiments: A Cahn-Hilliard analysis. *J. Chem. Phys.*, 117(13):6157–6168, 2002.

[4] J. F. Huang and L. S. Bartell. Kinetics of homogeneous nucleation in the freezing of large water clusters. *J. Phys. Chem.*, 99(12):3924–3931, 1995.

[5] S. V. Huffel and J. Vandewalle. *The Total Least Squares Problem: Computational Aspects and Analysis.* Society for Industrial and Applied Mathematics, 1991.

[6] L. Ickes, A. Welti, C. Hoose, and U. Lohmann. Classical nucleation theory of homogeneous freezing of water: Thermodynamic and kinetic parameters. *Phys. Chem. Chem. Phys.*, 17:5514–5537, 2015.

[7] D. Kashchiev. *Nucleation - Basic Theory with Applications.* Butterworth-Heinemann, 2000.

[8] T. S. Li, D. Donadio, G. Russo, and G. Galli. Homogeneous ice nucleation from supercooled water. *Phys. Chem. Chem. Phys.*, 13(44):19807–19813, 2011.

[9] A. Manka, H. Pathak, S. Tanimura, J. Wölk, R. Strey, and B. E. Wyslouzil. Freezing water in no-man's land. *Phys. Chem. Chem. Phys.*, 14(13):4505–4516, 2012.

[10] E. B. Moore and V. Molinero. Is it cubic? Ice crystallization from deeply supercooled water. *Phys. Chem. Chem. Phys.*, 13(44):20008–20016, 2011.

[11] B. J. Murray, S. L. Broadley, T. W. Wilson, S. J. Bull, R. H. Wills, H. K. Christenson, and E. J. Murray. Kinetics of the homogeneous freezing of water. *Phys. Chem. Chem. Phys.*, 12(35):10380–10387, 2010.

[12] B. J. Murray, D. O'Sullivan, J. D. Atkinson, and M. E. Webb. Ice nucleation by particles immersed in supercooled cloud droplets. *Chem. Soc. Rev.*, 41(19):6519–6554, 2012.

[13] T. Němec. Estimation of ice-water interfacial energy based on pressure-dependent formulation of classical nucleation theory. *Chem. Phys. Lett.*, 583:64–68, 2013.

[14] A. Pant, M. T. Parsons, and A. K. Bertram. Crystallization of aqueous ammonium sulfate particles internally mixed with soot and kaolinite: Crystallization relative humidities and nucleation rates. *J. Phys. Chem. A*, 110(28):8701–8709, 2006.

[15] M. T. Parsons, J. L. Riffell, and A. K. Bertram. Crystallization of aqueous inorganicmalonic acid particles: Nucleation rates, dependence on size, and dependence on the ammonium-to-sulfate ratio. *J. Phys. Chem. A*, 110(26):8108–8115, 2006.

[16] W. S. Price, H. Ide, and Y. Arata. Self-diffusion of supercooled water to 238 K using PGSE NMR diffusion measurements. *J. Phys. Chem. A*, 103(4):448–450, 1999.

[17] J. E. Shilling, M. A. Tolbert, O. B. Toon, E. J. Jensen, B. J. Murray, and A. K. Bertram. Measurements of the vapor pressure of cubic ice and their implications for atmospheric ice clouds. *Geophys. Res. Lett.*, 33(17):L17801, 2006.

[18] C. A. Stan, G. F. Schneider, S. S. Shevkoplyas, M. Hashimoto, M. Ibanescu, B. J. Wiley, and G. M. Whitesides. A microfluidic apparatus for the study of ice nucleation in supercooled water drops. *Lab Chip*, 9(16):2293–2305, 2009.